Complete Summary

TITLE

"Side Effects" scale of the Treatment Satisfaction Questionnaire for Medication (TSQM): the mean scale score for the "Side Effects" items on the TSQM (version II).

SOURCE(S)

Treatment Satisfaction Questionnaire for Medication (TSQM) (Version II). Quintiles Transnational Corp.; 2006. 3 p.

Measure Domain

PRIMARY MEASURE DOMAIN

Patient Experience

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the <u>Measure Validity</u> page.

SECONDARY MEASURE DOMAIN

Does not apply to this measure

Brief Abstract

DESCRIPTION

This "Side Effects" scale* on the Treatment Satisfaction Questionnaire for Medication (TSQM) is used to assess the level of dissatisfaction with the "Side Effects" of medication patients are taking. This composite scale is comprised of three of four items on the TSOM survey:

- As a result of taking this medication, do you experience any side effects at all? (y/n dichotomous skip)
- How dissatisfied are you by side effects that interfere with your physical health and ability to function (e.g., strength, energy levels)?
- How dissatisfied are you by side effects that interfere with your mental function (e.g., ability to think clearly, stay awake)?
- How dissatisfied are you by side effects that interfere with your mood or emotions (e.g., anxiety/fear, sadness, irritation/anger)?

*Note: The validity of this scale is influenced by the degree to which patients can; a) use personal experience to assess the effectiveness of their medication, b) has information on the clinical effectiveness of their medication (e.g., clinician appraisal, lab results etc.) and c) has formulated perceptions/beliefs about its effectiveness. Of note, patients' satisfaction and medication behavior may be the result of erroneous beliefs about the effectiveness of their medication. Moreover, medications may take time before they begin to work and TSQM assessments should be made after patients have had a chance to fully evaluate their medication.

Note: The "Side Effects" scale is a component of a composite measure. It cannot be used on its own and must be used with the following scales:

- "Effectiveness" scale of the Treatment Satisfaction Questionnaire for Medication (TSQM): the mean scale score for the "Effectiveness" items on the TSQM (version II).
- "Convenience" scale of the Treatment Satisfaction Questionnaire for Medication (TSQM): the mean scale score for the "Convenience" items on the TSOM (version II).
- "Global Satisfaction" scale of the Treatment Satisfaction Questionnaire for Medication (TSQM): the mean scale score for the "Global Satisfaction" items on the TSQM (version II).

RATIONALE

Patients' dissatisfaction with treatment may act as an early indicator of nonadherence and non-persistence with medication use, thereby affecting the clinical effectiveness and efficiency of medical care. Patients who perceive their medication to be ineffective (see the related National Quality Measures Clearinghouse (NQMC) summary, "Effectiveness" scale of the Treatment Satisfaction Questionnaire for Medication (TSQM): the mean scale score for the "Effectiveness" items on the TSOM (version II)), laden with side effects, very inconvenient (see the related NQMC summary, "Convenience" scale of the Treatment Satisfaction Questionnaire for Medication (TSQM): the mean scale score for the "Convenience" items on the TSQM (version II)), or as having more negative than positive characteristics (see the related NQMC summary, "Global Satisfaction" scale of the Treatment Satisfaction Questionnaire for Medication (TSQM): the mean scale score for the "Global Satisfaction" items on the TSQM (version II)) are less likely to either fill prescriptions or take their medications as prescribed. This in turn can impact the effectiveness of treatments and may result in service inefficiencies associated with treatment failure.

Treatment Satisfaction Questionnaire for Medication (TSQM) provides a unique opportunity to compare various medications used to treat a particular condition. Routine assessment of patients' level of TS-M provides a way for clinicians to screen individuals whose current medication experiences may increase the risk of poor medication adherence and weak persistence. If collected from many patients, such information can foster a deeper understanding of patients' perspectives when evaluating the merits and drawbacks of various treatment alternatives.

PRIMARY CLINICAL COMPONENT

Medication satisfaction; side effects

DENOMINATOR DESCRIPTION

Patients who completed the "Side Effects" items on the Treatment Satisfaction Questionnaire for Medication (TSQM)

NUMERATOR DESCRIPTION

The number of "Extremely Dissatisfied," "Very Dissatisfied," "Somewhat Dissatisfied," "Slightly Dissatisfied," "Not at all Dissatisfied," and "Not Applicable" responses for the "Side Effects" items on the Treatment Satisfaction Questionnaire for Medication (TSQM). From the responses, a scale score is calculated with a higher score indicating less dissatisfaction (greater satisfaction). A scale score is not computed for respondents indicating no side effects on the dichotomous question. For these individuals, the highest possible score on this scale may be assigned.

Evidence Supporting the Measure

EVIDENCE SUPPORTING THE CRITERION OF QUALITY

- Focus groups
- One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Unspecified

State of Use of the Measure

STATE OF USE

Current routine use

CURRENT USE

Internal quality improvement Quality of care research

Application of Measure in its Current Use

CARE SETTING

Ambulatory Care

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Nurses Pharmacists Physician Assistants Physicians Psychologists/Non-physician Behavioral Health Clinicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Individual Clinicians

TARGET POPULATION AGE

Age greater than or equal to 18 years

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

Unspecified

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

Unspecified

UTILIZATION

Unspecified

COSTS

Unspecified

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Getting Better Living with Illness Staying Healthy

IOM DOMAIN

Patient-centeredness

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

Patients who completed the "Side Effects" items on the Treatment Satisfaction Questionnaire for Medication (TSQM)

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

Patients who completed the "Side Effects" items on the Treatment Satisfaction Questionnaire for Medication (TSQM)

Exclusions

None

RELATIONSHIP OF DENOMINATOR TO NUMERATOR

All cases in the denominator are equally eligible to appear in the numerator

DENOMINATOR (INDEX) EVENT

Encounter

DENOMINATOR TIME WINDOW

Time window is a single point in time

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

The number of "Extremely Dissatisfied," "Very Dissatisfied," "Somewhat Dissatisfied," "Slightly Dissatisfied," "Not at all Dissatisfied," and "Not Applicable"

responses for the "Side Effects" items on the Treatment Satisfaction Questionnaire for Medication (TSQM). From the responses, a scale score is calculated with a higher score indicating less dissatisfaction (greater satisfaction). A scale score is not computed for respondents indicating no side effects on the dichotomous question. For these individuals, the highest possible score on this scale may be assigned.

Exclusions

None

MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

NUMERATOR TIME WINDOW

Encounter or point in time

DATA SOURCE

Patient survey

LEVEL OF DETERMINATION OF QUALITY

Not Individual Case

PRE-EXISTING INSTRUMENT USED

Unspecified

Computation of the Measure

SCORING

Non-weighted Score/Composite/Scale

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

External comparison at a point in time External comparison of time trends Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

A multi-step exploratory factor analysis (EFA) was employed to investigate the construct validity of the Treatment Satisfaction Questionnaire for Medication (TSQM). The TSQM v. II was tested for model fit against an established theoretical model (the Decisional Balance Model of Treatment Satisfaction) using hierarchical confirmatory factor analysis (HCFA).

EVIDENCE FOR RELIABILITY/VALIDITY TESTING

Atkinson MJ, Kumar R, Cappelleri JC, Hass SL. Hierarchical construct validity of the treatment satisfaction questionnaire for medication (TSQM version II) among outpatient pharmacy consumers. Value Health2005 Nov-Dec;8 Suppl 1:S9-S24. PubMed

Atkinson MJ, Sinha A, Hass SL, Colman SS, Kumar RN, Brod M, Rowland CR. Validation of a general measure of treatment satisfaction, the Treatment Satisfaction Questionnaire for Medication (TSQM), using a national panel study of chronic disease. Health Qual Life Outcomes2004 Feb 26;2:12. [77 references]

PubMed

Bharmal M, Payne K, Atkinson MJ, Desrosiers MP, Morisky DE, Gemmen E. Validation of an abbreviated Treatment Satisfaction Questionnaire for Medication (TSQM-9) among patients on antihypertensive medications. Health Qual Life Outcomes2009;7:36. PubMed

Identifying Information

ORIGINAL TITLE

Side Effects.

COMPOSITE MEASURE NAME

Treatment Satisfaction Questionnaire for Medication (TSQM) Version II

SUBMITTER

Atkinson, Mark J, Ph.D.

DEVELOPER

Atkinson, Mark J, Ph.D.; Hass, Steven L, Ph.D.

FUNDING SOURCE(S)

Pharmacia, Pfizer Inc.

COMPOSITION OF THE GROUP THAT DEVELOPED THE MEASURE

Atkinson, Mark J, Ph.D.; Hass, Steven L, Ph.D.; Kumar, Ritesh, Ph.D.; Colman, Shoshana, Ph.D.; Sinha, Anusha, M.P.H.; Brod, Meryl, Ph.D.; Roland, Clayton, Ph.D.

FINANCIAL DISCLOSURES/OTHER POTENTIAL CONFLICTS OF INTEREST

Unspecified

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2006 Jan

MEASURE STATUS

This is the current release of the measure.

SOURCE(S)

Treatment Satisfaction Questionnaire for Medication (TSQM) (Version II). Quintiles Transnational Corp.; 2006. 3 p.

MEASURE AVAILABILITY

The individual measure, "Side Effects," is published in "Treatment Satisfaction Questionnaire for Medication (TSQM)." This document is available from the Quintiles Web site.

For more information contact, Quintiles, Inc., 3130 Fairview Park Drive, Suite #501, Falls Church, VA 22042; Telephone: 703-269-5857; Fax: 703-269-5802; Web site: www.quintiles.com.

NQMC STATUS

This NQMC summary was completed by ECRI Institute on September 8, 2009. The information was verified by the measure developer on October 15, 2009.

COPYRIGHT STATEMENT

Quintiles, Inc. holds the copyright for the Treatment Satisfaction Questionnaire for Medication (TSQM). Any uses of the TSQM must obtain permission for use from

Quintiles. There is a per-use fee associated with the TSQM. If you are interested in using the TSQM, please contact: Murtuza Bharmal, BPharm, Ph.D., Analytics & Health Outcomes, Quintiles, Inc., TSQM@Quintiles.com or murtuza.bharmal@Quintiles.com.

Copyright/Permission Requests

Date Modified: 1/4/2010

